

Taxes on sugary drinks: Why do it?




Sugary drinks¹ – a major contributor to obesity and diabetes

- Over-consumption of sugar is a major contributor to obesity, diabetes and tooth decay.
- In the current food environment it is very easy to consume too much sugar, especially from sugary drinks.
- Sugary drinks are a major source of sugar in the diet, and its consumption is increasing in most countries, especially amongst children and adolescents.
- On average, a single can of a sugary drink contains around 40 grams of free sugars² (equivalent to around 10 teaspoons of table sugar).
- WHO guidelines recommend that, to prevent obesity and tooth decay, adults and children reduce their consumption of free sugars to less than 10% of their daily energy intake (equivalent to around 12 teaspoons of table sugar for adults). The guidelines suggest further reducing intake of sugars to below 5% of daily energy intake (around 6 teaspoons of table sugar for adults) for additional health benefits (1).

¹ Sugary drinks are products that contain added sugar, corn or fruit-juice concentrates and include carbonates, fruit drinks, sports drinks, energy and vitamin water drinks, sweetened iced tea, and lemonade.

² Free sugars refer to monosaccharides (such as glucose, fructose) and disaccharides (such as sucrose or table sugar) added to foods and drinks by the manufacturer, cook or consumer, and sugars naturally present in honey, syrups, fruit juices and fruit juice concentrates

Percentage of adolescents who drink soft drinks daily



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

Data Sources: Global School-based Student Health Survey; Health Behaviour in School-aged Children (HBSC) Study; International Report from the 2009/2010 Survey.

Map Production: Information Evidence and Research (IER) World Health Organization



© WHO 2016. All rights reserved.

Problem

- The worldwide prevalence of obesity more than doubled between 1980 and 2014.
- An estimated 39% of adults were overweight in 2014, and 13% were obese (2).
- Some 42 million children under the age of 5 were overweight or obese in 2015 (3).
- The prevalence of overweight in pre-school aged children is increasing fastest in low- and lower middle-income countries.
- People who consume sugary drinks regularly – 1 to 2 cans a day or more – have a 26% greater risk of developing type 2 diabetes than people who rarely consume such drinks (4).
- The number of people with diabetes has risen from 108 million in 1980 to 422 million in 2014 (5).
- Apart from diabetes, obesity is a major risk factor for heart diseases, cancers and other diseases.

Costs

- From 2011 to 2030, losses in gross domestic product worldwide due to diabetes, including both direct and indirect costs, are expected to total US\$ 1.7 trillion, US\$ 900 billion in high-income countries and US\$ 800 billion in low- and middle-income countries (6).

What we can do

Governments can take a number of actions to improve availability and access to healthy foods and have a positive influence on the food people choose to consume. A major action for comprehensive programmes aimed at reducing consumption of sugars is taxation of sugary drinks. Just as taxing tobacco helps to reduce tobacco use, taxing sugary drinks can help reduce consumption of sugars.



Benefits

Taxes on sugary drinks help reduce consumption and prevent obesity

- Evidence shows that a tax of 20% on sugary drinks can lead to a reduction in consumption of around 20%, thus preventing obesity and diabetes (7).

Savings on healthcare

- Estimates suggest that, over 10 years, a tax on sugary drinks of 1 cent per ounce in the United States of America would result in more than US\$ 17 billion in healthcare cost savings (8).

Revenues raised from taxes can be used to promote the health of the population

- This tax could generate approximately US\$ 13 billion in annual tax revenues in the United States of America in 2016 (9).

- Based on 2014 data, a tax on sugary drinks of 1 yuan (US\$ 0.16) per litre in China would generate an estimated 73.6 billion yuan (US\$ 11.8 billion) in revenues (10).
- Revenue generated by these taxes could be spent on efforts to improve health care systems, encourage healthier diets, increase physical activity, or build capacity for effective tax administration, further increasing the value of this measure.

Low-income consumers and young people get the greatest health benefits from taxes

- In Mexico, a year after the introduction of a tax on sugary drinks, households with the fewest resources reduced their purchases of sugary drinks by 17%, compared to 12% for the general population (11).

To reduce over-consumption of sugars and halt the epidemic of obesity and diabetes, countries need comprehensive action plans that combine taxation, restriction of marketing of sugary products to children, and education.

CASE STUDY

Evidence shows that implementing taxes on sugary drinks leads to reduced consumption of these products. Several countries are well on their way to implementing taxes on sugary drinks.

In January 2014, the government of Mexico added a 1 peso per litre excise tax on any non-alcoholic beverage with added sugar (powder, concentrates or ready-to-drink) to the country's Special Tax on Production and Services, which is paid by the producer and represents about a 10% increase in price for the consumer.

A study conducted by the Mexican National Institute of Public Health and the University of North Carolina evaluating the first year of implementation showed

an average reduction of 6% in the purchase of taxed sugary drinks during 2014. This reduction increased over the course of the year to reach 12% by December 2014. Households with the fewest resources had an average reduction in purchases of 9% in 2014, increasing to 17% by December. The study showed a 4% increase in purchases of untaxed beverages, particularly purchased bottled water (12).

Over US\$ 2.6 billion was raised during the first two years of implementation; some of this revenue is beginning to be invested towards installing water fountains in schools across Mexico (13).



References

1. Guideline: Sugars intake for adults and children. Geneva: World Health Organization; 2015 (http://apps.who.int/iris/bitstream/10665/149782/1/9789241549028_eng.pdf).
2. Global status report on noncommunicable diseases 2014. Geneva: WHO; 2014.
3. Levels and Trends in Child Malnutrition. UNICEF/WHO/World Bank Group Joint Child Malnutrition Estimates. Key findings of the 2016 edition. New York: UNICEF, Geneva: WHO, Washington DC: World Bank Group; 2016.
4. Malik VS, Popkin BM, Bray GA, Despres JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care*, 2010;33:2477-83.
5. Global report on diabetes. World Health Organization: Geneva; 2016.
6. Bloom DE, Cafiero ET, Jané-Llopis E, Abrahams-Gessel S, Bloom LR, Fathima S, et al. The global economic burden of noncommunicable diseases (Working Paper Series). Geneva: Harvard School of Public Health and World Economic Forum; 2011.
7. Powell, LM., Chriqui JF, Khan T, Wada R, Chaloupka FJ. Assessing the potential effectiveness of food and beverage taxes and subsidies for improving public health: a systematic review of prices, demand and body weight outcomes. *Obesity Reviews*, 2013; 14:110-128.
8. Wang YC, Coxson P, Shen Y, Goldman L, Bibbins-Domingo K. A penny-per-ounce tax on sugar-sweetened beverages would cut health and cost burdens of diabetes. *Health Affairs*, 2014; 31, no 1: 199-207.
9. Rudd Center for Food Policy & Obesity. Revenue Calculator for Sugar-Sweetened Beverage Taxes. Available at: www.uconnruddcenter.org/revenue-calculator-for-sugar-sweetened-beverage-taxes.
10. Chaloupka FJ. Existing Evidence and Guidance on Fiscal Policies. Presentation in Technical Meeting on Fiscal Policies for Diet and Prevention of Noncommunicable Diseases, 5-6 May 2015, Geneva, Switzerland.
11. Colchero MA, Popkin BM, Rivera JA, Ng SW. Beverage purchases from stores in Mexico under the excise tax on sugar sweetened beverages: observational study. *BMJ*, 2016 Jan 6;352:h6704.
12. Idem.
13. Fact Sheet. Uncapping the Truth: The Mexican Sugar Sweetened Beverage Tax Works. The Nutritional Health Alliance: Mexico; 2016.



For more information,
please contact the Department
of Prevention of Noncommunicable
Diseases, WHO, at hq_pnd@who.int
(www.who.int/dietphysicalactivity/en)

#Tax4Health



© World Health Organization 2016.

All rights reserved.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the materials lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

WHO/NMH/PND/16.5